



## Preparation for Your Upcoming Colonoscopy

### YOU WILL NEED:

- Plenvu® prescription filled by your pharmacy

If you didn't collect your prep prescription shortly after we sent it to your pharmacy, kindly contact your pharmacy to have it prepared again.

### Appointment Info

DATE \_\_\_\_\_

ARRIVAL TIME \_\_\_\_\_

- ☐ **Endoscopy Center of Marin**  
1100 S. Eliseo Dr. Suite 3, Greenbrae  
Expect to be at the center 90 minutes
- ☐ **MarinHealth Medical Center**  
250 Bon Air Road, Greenbrae  
Expect to be at the hospital 2 hours
- ☐ **Novato Community Hospital**  
180 Rowland Way, Novato  
Expect to be at the hospital 3 hours

### Diabetes Instructions (if applicable)

Oral Medication Instructions  
\_\_\_\_\_  
Insulin Schedule  
\_\_\_\_\_

### Blood Thinner Instructions (if applicable)

- ☐ Continue blood thinner as prescribed
- ☐ HOLD for \_\_\_\_\_ days

### Weight Loss Prescriptions (if applicable)

- ☐ Weekly Dosing(Ozempic, Rybelsus, Wegovy, Trulicity, Mounjaro) Your last dose must be done 7 days before your procedure.
- ☐ Daily Dosing(Saxenda, Byetta) Do not take on the day of the procedure.
- ☐ Other instruction:  
\_\_\_\_\_

### MarinHealth Gastroenterology | A UCSF Health Clinic

200 Tamal Plaza | Suite 200 | Corte Madera, CA 94925

7120 Redwood Blvd. Novato, CA 94945

415-925-6900 | MyMarinHealth.org/gastroclinic





## 7 Days Before Your Colonoscopy

- Confirm ride for the day of your procedure
- If you take blood thinners or weight loss medication, please **review instructions** for holding prescription.

**Last chance to cancel/reschedule your procedure appointment. Please be aware that the Endoscopy Center of Marin may impose a late cancellation charge of up to \$200.**



## 5 Days Before Your Colonoscopy

- Review **thorough prep instructions** and plan your meals
- Create a list of your medications and supplements to provide the facility. Include prescription name, dose, and directions
- Create a list of your allergies to provide the facility



## 3 Days Before Your Colonoscopy

- STOP eating foods with nuts, seeds, or whole grains. (fruits and vegetables with seeds, sesame, etc.)
- Avoid corn, beans, quinoa, and popcorn
- For more diet recommendations, **please visit:** [MyMarinHealth.org/prep](http://MyMarinHealth.org/prep)



## 2 Days Before Your Colonoscopy

- Increase your fluid intake, drink a minimum of 64 oz of clear liquids (approximately 8 glasses) (water preferred)
- Eat well-balanced meals, but continue avoiding nuts, seeds, and whole grains



## 1 Day Before Your Colonoscopy

### “PREP DAY”

- Have a low-residue breakfast BEFORE 10 am (white toast, eggs, yogurt, cream of wheat, smoothies without seeds)
- At 10 am begin a “Clear Liquid Diet.” Clear liquids include water, juice without pulp, soda, black coffee/tea (no milk), electrolyte drinks, Ensure CLEAR, Boost Breeze®, Jell-O, broth, popsicles
- At 6 pm drink the first dose of Plenvu®. Empty Dose 1 into mixing container. Add water to fill line (16 oz line) and mix until completely dissolved. Drink the full amount within 30 minutes
- After drinking the Plenvu® mix, drink 16 oz of clear liquids within 30 minutes
- Continue drinking clear liquids until bedtime

## The Day of Your Colonoscopy

- Wake up 5 – 6 hours before your arrival time to start the **second** dose. Then drink the second dose of Plenvu®
- Empty dose 2 (Pouch A and Pouch B) into mixing container. Add water to fill line (16 oz line) and mix until dissolved. Drink full amount within 30 minutes
- **Please complete the bowel prep 3 hours before your colonoscopy arrival time. Please STOP drinking all liquids 3 hours before your colonoscopy arrival time.**
- You should take your regular medications/ supplements at least three hours before your arrival time or after your colonoscopy.

### BRING THE FOLLOWING:

- Photo ID
- Insurance Card
- Medication List
- Allergy List

## TIPS

- To improve the taste try chilling laxative and/or using a straw to drink mixture
- If you experience nausea or vomiting, rest for 15 minutes, then try to resume prep
- Use unscented wipes instead of toilet paper.
- Use Vaseline or an ointment for diaper rash around the anus before and/or during prep to minimize irritation from passing bowel movements